



## **GCN CIRCULAR 20366, LIGO/Virgo G268556: INTEGRAL search of temporally coincident prompt hard X-ray emission**

**Savchenko, V.; Ferrigno, C.; Mereghetti, S.; Kuulkers, E.; Bazzano, A.; Bozzo, E.; Courvoisier, T. J.L.; Brandt, Søren; Diehl, R.; Hanlon, L.**

*Total number of authors:*  
15

*Publication date:*  
2017

*Document Version*  
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

*Citation (APA):*  
Savchenko, V., Ferrigno, C., Mereghetti, S., Kuulkers, E., Bazzano, A., Bozzo, E., Courvoisier, T. J. L., Brandt, S., Diehl, R., Hanlon, L., Laurent, P., Lutovinov, A., Roques, J. P., Sunyaev, R., & Ubertini, P. (2017, Jan 4). GCN CIRCULAR 20366, LIGO/Virgo G268556: INTEGRAL search of temporally coincident prompt hard X-ray emission. <https://gcn.gsfc.nasa.gov/gcn3/20366.gcn3>

---

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

TITLE: GCN CIRCULAR  
 NUMBER: 20366  
 SUBJECT: LIGO/Virgo G268556: INTEGRAL search of temporally coincident prompt hard X-ray emission  
 DATE: 17/01/04 22:55:59 GMT  
 FROM: Carlo Ferriño at ISDC/INTEGRAL <carlo.ferriño@uni.ge.ch>

V. Savchenko and C. Ferriño (ISDC, University of Geneva, CH)  
 on behalf of the INTEGRAL group: S. Mereghetti (IASF-Mi, Italy),  
 E. Kuulkers (ESTEC/ESA, The Netherlands),  
 A. Bazzano (IAPS-Roma, Italy),  
 E. Bozzo, T. J. -L. Courvoisier (ISDC, University of Geneva, CH)  
 S. Brandt (DTU - Denmark)  
 R. Diehl (MPE-Garching, Germany)  
 L. Hanlon (UCD, Ireland)  
 P. Laurent (APC-CeA, France)  
 A. Lutovinov (IKI, Russia) J. P. Roques (CESR, France)  
 R. Sunyaev (IKI, Russia)  
 P. Ubertini (IAPS-Roma, Italy)

We investigated serendipitous INTEGRAL observations carried out at the time of the LIGO/Virgo G268556. The satellite was pointing at RA = 00:04:02 Dec=+67:14:38, away from the high-probability region, derived from the LIGO Bayestar pipeline. The anti-coincidence shield of spectrometer on board of INTEGRAL (SPI/ACS) covered the full LIGO 90% confidence region and provided

the most stringent constraints on the flux of a possible electromagnetic counterparts in the energy range covered by INTEGRAL instruments. We investigated the SPI-ACS light curve between -100 and +100 s from the trigger time (2017-01-04 10:11:59 UTC) on temporal scales from 0.1 to 10 s and found no significant excesses over a very stable background.

The SPI/ACS light curves, binned at 50 ms, are derived from 91 independent detectors with different lower energy thresholds (mainly between 50 keV and 150 keV) and an upper threshold at about 100 MeV.

The ACS response varies as a function of the incident angle. Assuming an optimal perpendicular direction of the burst to the INTEGRAL pointing direction, we estimate a 3-sigma upper limit corresponding to fluences of  $1.6 \times 10^{-7}$  erg/cm<sup>2</sup> for a 1 s duration,  $4.5 \times 10^{-7}$  erg/cm<sup>2</sup> for 10 s, and  $5 \times 10^{-8}$  erg/cm<sup>2</sup> for 0.1 s. For this computation, we adopt a low threshold at 100 keV and Band model parameters  $\alpha' 0.5$ ,  $\alpha' 2.5$  with peak  $E_0 \sim 600$  keV.

The optimal orientation is compatible with a large part of the high-probability sky region of the trigger. The limit anywhere in the 90% confidence range of the LIGO Bayestar localisation is at most 50% less stringent.

Investigations of the light curves from the other INTEGRAL instruments covering part of the LIGO localisation region (IBIS VETO, ISGRI, and PICsIT) did not reveal any significant transient event in the 200 s temporal window centred on LIGO/Virgo G268556, but provide less stringent upper limit on the fluence.